

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/98323/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Yeo, Syn Kok, French, Rhiannon, Filomena, Spada and Clarkson, Richard
ORCID: <https://orcid.org/0000-0001-7389-8673> 2017. Opposing roles of Nfkb2 gene products p100 and p52 in the regulation of breast cancer stem cells. Breast Cancer Research and Treatment 162 (3) , pp. 465-477. 10.1007/s10549-017-4149-0 file

Publishers page: <http://dx.doi.org/10.1007/s10549-017-4149-0>
<<http://dx.doi.org/10.1007/s10549-017-4149-0>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



[Click here to view linked References](#)

Opposing roles of *Nfkb2* gene products p100 and p52 in the regulation of breast cancer stem cells

Syn Kok Yeo^{1*}, Rhiannon French², Filomena Spada² and Richard Clarkson²

¹Department of Cancer Biology, University of Cincinnati, Vontz Building, 3125 Eden Avenue, Cincinnati Ohio 45267.

²European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, Hadyn Ellis Building, Maindy Road, Cathays, Cardiff, CF24 4HQ

*Corresponding author: Syn Kok Yeo

Department of Cancer Biology, University of Cincinnati, Vontz Building, 3125 Eden Avenue, Cincinnati Ohio 45267.

Phone: 513-558-4560

Fax: 513-558-5050

Email: yeosn@ucmail.uc.edu

Abstract

Purpose: Nuclear factor-kappa B (NF-κB) signalling has been shown to regulate properties of breast cancer stem cells. However, the specific contribution of the non-canonical NF-κB pathway, components of which are elevated in aggressive breast cancer has not been addressed.

Methods: Through shRNA silencing of the Nfkb2 gene, the role of p100/p52 in 4T1 and N202.1A cell lines were assessed by NF-κB reporter, invasion, tumoursphere and orthotopic transplantation assays. The processing of p100 into p52 was also inhibited with a p97 ATPase inhibitor, NMS-873, and its effects on tumoursphere formation was assessed.

Results: Knockdown of Nfkb2 led to opposing changes in NF-κB dependent transcription. NF-κB activity was elevated in 4T1 cells and this resulted in increased motility, cancer stem cell (CSC) activity and tumourigenicity in vivo. Conversely, depletion of Nfkb2 in N202.1a cells decreased NF-κB activity, CSC properties and tumourigenicity in vivo. By selectively over-expressing the p52 subunit in Nfkb2 depleted cells, we found that the increased malignancy in 4T1 cells could not be reverted in the presence of p52 whereas the decreased tumourigenicity of N202.1a cells could be rescued by p52. These results indicate that p100 and its subunit p52 have opposing effects on breast CSC activity. Accordingly, inhibition of an upstream regulator of p100 processing was effective in reducing tumoursphere formation of N202.1A and SKBR3 (ErbB2^{HIGH}) cells without aggravating that of 4T1 and MDA-MB-231 (ErbB2^{LOW}) cells.

Conclusion: These findings indicate that inhibiting the processing of p100 may be a potential therapeutic strategy to suppress CSC activity in a subset of breast tumours.

Keywords: breast cancer, Nfkb2, NF-KappaB, cancer stem cell, ErbB2, p97 ATPase

Introduction

Even though breast cancer predominantly affects women, it is the second most common form of cancer globally [1-4]. Significant advances have led to improved survival rates but the reality is that breast cancer is a heterogeneous disease which comprises of varying subtypes [5, 6]. While prognoses have improved for certain subtypes, some remain less adequate. This is mainly due to therapeutic resistance, metastasis and tumour relapse, which can be attributed at least in part to a subpopulation of cells with increased tumor initiating potential, termed cancer stem cells.

The existence of cancer stem cells (CSCs) were first shown in hematopoietic malignancies [7, 8] and more recently, the presence of CSCs in breast tumours represented the first indication of CSCs in solid tumours [9]. The CSC hypothesis states that cells within a tumour are not equal in their tumorigenic potential. Evidence suggests that within this hierarchy, it is the CSCs that fuel the growth of a tumour resulting in therapeutic resistance, disease relapse and the seeding of metastases in a range of cancer subtypes [10, 11]. CSCs have been shown to utilize and depend on distinct signalling pathways from non-CSCs and this suggests a possibility for specific therapeutic targeting to eradicate CSCs [12-14].

One such candidate pathway involves NF- κ B dependent transcription. NF- κ B signalling has been shown to promote breast CSC properties [35] and to result in EMT and metastasis in animal models of breast cancer [28-30]. The NF- κ B proteins are transcription factors which were initially identified as regulators of the immune system. There are five different subunits which can form homo- or heterodimers to regulate the expression of genes. Namely, these are RelA (p65), RelB, c-rel, p50 and p52. All of these subunits have a conserved Rel homology domain

1
2
3
4 which facilitates DNA binding and dimerization. However, p50 and p52 differ from the Rel
5
6 proteins in two ways. Firstly, they lack trans-activating domains which are important for
7
8 initiating transcription. Secondly, p50 and p52 are translated as precursor proteins p105 and p100
9
10 respectively. These precursor proteins are inhibitor of NF- κ B (I κ B) proteins characterized by the
11
12 presence of ankyrin repeats at the C-terminus. These motifs are removed upon partial
13
14 proteasomal processing into p50 or p52 respectively. Apart from p105 and p100, other I κ B
15
16 proteins include I κ B α , I κ B β , I κ B ϵ and Bcl-3. The I κ B proteins function by masking the nuclear
17
18 localization signal of NF- κ B subunits and thus retain them in the cytoplasm. There are two main
19
20 NF- κ B pathways, namely, the canonical and alternative pathways. Transcriptional activity
21
22 induced by canonical NF- κ B signalling involves RelA-p50 dimers whereas the alternative
23
24 pathway is dependent on RelB-p52 dimers. The alternative pathway, which is associated with
25
26 constitutive NF- κ B activity in normal and tumour cells is maintained through a signalling
27
28 cascade involving NF- κ B inducing kinase (NIK) and I κ K α dimers, resulting in p100
29
30 phosphorylation and processing into p52. The loss of ankyrin domains from the C-terminus of
31
32 p100 then allows translocation of RelB-p52 heterodimers into the nucleus [15, 16]. P100 also
33
34 actively shuttles between the nuclear and cytoplasm and plays an inhibitory role by sequestering
35
36 other NF- κ B subunits in the nucleus [17, 18]. The processing of p100 into p52 has also been
37
38 recently shown to depend on the ATPase activity of p97 and this presents a critical regulation
39
40 point where the alternative pathway can be inhibited [19]. Taken together, the roles of the Nfkb2
41
42 gene which encodes both the p100 inhibitory subunit and p52 transcription factor is likely to be
43
44 complex and context dependent.
45
46
47
48
49
50
51
52
53
54
55

56 As breast cancers are scrutinized further into more distinct subtypes [20], it is likely that
57
58 more specific therapies will lead to more effective targeting of each particular subtype with less
59
60
61
62
63
64
65

side effects. The canonical NF- κ B pathway has been shown to be important for tumorigenesis and CSC properties especially in estrogen receptor (ER) negative and ErbB2 driven tumours [21-24], yet the alternative pathway is also known to play a key role in driving oncogenic pathways such as inducing cyclin D1 expression [25, 26]. Aberrant levels of alternative NF- κ B pathway components in human breast cancer tissue have been described [27, 28]. However, since p100 and p52 are known to play opposing roles in the regulation of the alternative NF- κ B pathway, we set out to determine whether certain breast cancers may respond differentially to silencing of the *Nfkb2* gene, and whether this acts on breast CSC viability. Interestingly, we found that depletion of *Nfkb2* has context dependent effects on 4T1 (ErbB2^{low}) and N202.1A (ErbB2^{high}) mammary cancer cell lines respectively. NF- κ B activity was found to be differentially affected in these cell lines upon *Nfkb2* silencing and this led to corresponding phenotypes associated with cancer stem cell properties and activity. While our results corroborate previous findings where *Nfkb2*-mediated transcription was shown to be important for breast cancer stem cell activity [29], we provide evidence that selective targeting of p52 rather than upstream *Nfkb2*/p100 expression may have a more beneficial therapeutic potential in the in vivo setting.

Materials and Methods

Reagents and antibodies

Plasmid containing human p52 sequence was a kind gift from Professor Neil Perkins (Newcastle University). This HA-tagged p52 sequence was excised by XbaI-PstI digestion and ligated into the cloning site of P305 lentiviral plasmid. (Dr. Riccardo Brambilla, Cardiff University). NF- κ B luciferase assays were carried out using the 3x κ B luciferase reporter plasmid (Professor Ron Hay, University of St. Andrews) and pcDNA3.1-LacZ plasmid (Invitrogen) was used as control for transduction efficiency. Packaged lentiviral vectors encoding shRNA against *Nfkb2* (NM_019408) or non-target (NT) (SHC002V) control sequences were obtained from Sigma. Antibodies for Western blotting against β -actin (Abcam, Ab8227), Lamin A/C (CST 4777), p65 (CST 8243), p105/p50 (eBioscience 14-6732-81) and p100/p52 (SantaCruz, SC-7386) were used according to manufacturer's instructions. For flow cytometry experiments, CD29 (BD Biosciences, 562155), CD61 (eBioscience, 13-0611-81), CD44 (Biolegend, 103007) and CD24 (Biolegend, 311106) antibodies were used.

Cell culture

4T1 cells [30] were obtained from Dr. Robin Anderson (University of Melbourne) and N202.1A cells [31] were a kind gift from Dr. Pier-Luigi Lollini (Sezione di Cancerologia, Italy). 4T1 and SKBR3 cells were cultured in RPMI medium with 10% fetal bovine serum (FBS). N202.1A and MDA-MB-231 cells were cultured in DMEM medium that is supplemented with FBS at 10%. All media were also supplemented with 2mM L-glutamine and 50units/ml penicillin-streptomycin. For generation of stable cell lines, cells transduced with respective lentivirus were then cultured in 4 μ g/ml puromycin for 4T1 cells and 1 μ g/ml puromycin for

1
2
3
4 N202.1a. For BAY 11-7082 (Cayman Chemical) and NMS-873 (Apexbio) experiments,
5
6
7 mammary cancer cells were maintained with 1 μ M BAY 11-7082 or NMS-873 throughout the
8
9 time period of the assay.
10

11 12 **NF- κ B luciferase reporter assays** 13

14
15 Cells were seeded in clear bottom black well plates and transfected 24 hours later with
16
17 NF- κ B reporter plasmid along with pcDNA3.1-LacZ as control for transfection efficiency using
18
19 Lipofectamine 2000 (Invitrogen). After 48 hours, cells were lysed with Glo-lysis buffer
20
21 (Promega). Luciferase and LacZ activity was measured after adding Bright-Glo (Promega) and
22
23 Beta-Glo (Promega) substrates according to the manufacturer's instructions. Resulting
24
25 luminescence were measured using a Flurostar Optima plate reader (BMG Labtech).
26
27
28
29
30

31 **Transwell migration assay** 32

33
34 Cells were dissociated into single cell suspension and seeded into each 24-well cell
35
36 culture insert with 8 μ m pores (BD Biosciences). Cells were seeded with respective medium
37
38 containing 0.1% FBS at a density of 25,000 cells/chamber. Migration was stimulated by a serum
39
40 gradient. After 24 hours, cells were fixed with ice cold 70% ethanol. Cells on the upper surface
41
42 were then removed mechanically and the membranes were stained with haematoxylin and eosin
43
44 for visualization.
45
46
47
48
49

50 **Mammosphere forming assay** 51

52
53 Cells were dissociated into single cell suspensions and re-suspended in mammosphere
54
55 culture medium (MEBM, Lonza) supplemented with B27, 5mg/ml insulin, 20ng/ml Epidermal
56
57 growth factor, 1mg/ml hydrocortisone and 0.0008% v/v β -mercapthoethanol. Cells were seeded
58
59
60
61
62
63
64
65

at a density of 500cells/well in ultra-low attachment 96-well plates (Corning) and incubated for 7 days. Passaging was achieved by dissociating mammospheres into single cell suspensions and replating under similar conditions.

qRT-PCR analysis

Isolation of RNA from cells was carried out using the RNeasy Mini Kit (Qiagen, Sussex, UK) according to manufacturer's instructions. Then, cDNA was synthesized from RNA samples using RevertAID Premium Reverse Transcriptase (Fermentas). SYBR Green reagent (Invitrogen) was used in the reaction mastermix and thermal cycling was carried out with a Step One Plus Realtime PCR System (Applied Biosystems). Cycling conditions used for all reactions consisted of; initial denaturation for 10 minutes at 95°C, followed by 45 cycles of 95°C for 15 seconds and 60°C for 1 minute. List of primers used are: GAPDH F-CACAGTCAAGGCCGAGAATG, GAPDH R-TCTCGTGTTCACACCCATC, SNAIL F-TGAGAAGCCATTCTCCTGCT, SNAIL R-CTTCACATCCGAGTGGGTTT, SLUG F-TCTGCAGACCCACTCTGATG, SLUG R-AGCAGCCAGACTCCTCATGT, TWIST1 F-CAGGCCGGAGACCTAGATGTCAT, TWIST1 R-TGCCCCACGCCCTGATTCTTG, TWIST2 F-CTTCTCCGTGTGGCGCAT, TWIST2 R-CCGCATCCAGTTCCGCAT, ZEB2 F-CCGGGAGCTGTTTCTTCG, ZEB2 R-CGCAGGCTCGATCTGTGA, ECADHERIN F-CAGATGATGATACCCGGGACAA, ECADHERIN R-GGAGCCACATCATTTCGAGTCA, NCADHERIN F-AGGACCCTTTCCTCAAGAGC, NCADHERIN R-ATAATGAAGATGCCCGTTGG. C_T values obtained were processed using the 2^{-ΔΔC_T} relative quantification method [32].

Western Blotting

Protein lysates were collected from cultured cells by lysing with RIPA buffer (150mM sodium chloride, 1% v/v Nonidet-P40, 0.1% w/v sodium dodecyl sulphate, 0.5% w/v sodium deoxycholate, 50mM Tris pH8) with the addition of complete mini protease inhibitor tablets (Roche) and phosphatase inhibitors (10mM sodium fluoride, 1mM sodium orthovanadate, 10mM sodium pyrophosphate). Proteins were then separated by SDS-PAGE and immune-blotted for detection of proteins.

Flow cytometry analysis

Cells were reconstituted in single cell suspensions and incubated with antibodies for 20 minutes at 4°C. Flow analysis was carried out on a BD FACSCanto cell analyzer.

Orthotopic transplantation of mammary cancer cells and tumour measurements

For transplantation experiments, BalbC and NOD/SCID-BalbC mice of six to eight weeks of age were obtained from Charles River Laboratories. Maintenance, breeding and scientific procedures involving animals were carried out according to the guidelines set by the U.K. Home Office Regulations Animals (Scientific Procedures) Act 1986. Orthotopic cell transplants were carried out on mice between 8-12 weeks of age. For 4T1 transplants, cells were dissociated into single cell suspension and 10,000 cells were injected into the abdominal mammary glands. For N202.1a transplants, 50,000 cells were injected into NOD/SCID-BalbC mice.

Statistical analyses

Data are presented as means \pm SEM. For statistical analyses, two-tailed unpaired student's t-test was used and the threshold for significance was $p < 0.05$. For tumor growth curves, ANCOVA test was used to determine significance.

Results

Knockdown of *Nfkb2* led to elevated basal NF- κ B activity in 4T1 cells but decreased basal NF- κ B activity in N202.1a cells

Through the analysis of The Cancer Genome Atlas (TCGA) breast datasets, we found elevated expression of NFKB2 transcripts in invasive breast carcinomas relative to normal breast tissue (Fig. 1a). This suggests a role for NFKB2 dependent signalling in advanced breast cancer. Consequently, we went on to address the direct contribution of *Nfkb2* to the tumourigenic properties of two invasive mammary cancer cell lines, N202.1A and 4T1, by shRNA knockdown. NF- κ B activity has previously been reported to be associated with ErbB2 levels in breast cancer [26,27], and so as expected, the N202.1A cell line which was derived from an ErbB2 (Neu) over-expressing mammary tumour had concomitantly elevated nuclear p65 levels determined by western blot analysis compared to the ErbB2^{LOW} mammary cell line 4T1 (Figure 1b). In contrast however, the relative levels of nuclear p100/p52 and the ratio of p52 to p100 was higher in 4T1 cells relative to N202.1A cells. After silencing *Nfkb2*, the nuclear p52 to p100 ratio, as well as the nuclear to cytoplasmic p52 ratio, were further increased in 4T1 cells but decreased in N202.1A cells (Fig. 1b-1d) while no significant changes in the levels of p105, p50 or p65 were observed in both nuclear and cytoplasmic fractions upon *Nfkb2* knockdown. These relative changes in p52/p100 correlated with basal NF- κ B activity as indicated by NF- κ B reporter assay (Fig 1e). The different outcomes in activation of NF- κ B in the cell lines examined could be due to varying levels of upstream signals which affect basal processing of p100 into p52. Irrespective of the underlying mechanism for this difference however, it is clear that perturbation of p100/p52 levels affect basal NF- κ B activity and this demonstrates the contribution of non-canonical NF- κ B pathway proteins towards overall basal NF- κ B activity in

1
2
3
4 these mammary tumour cell lines. Given the presumptive role for NF- κ B signalling in tumour
5
6 growth and malignancy, we set out to evaluate whether the differential effects on NF- κ B activity
7
8 mediated by *Nfkb2* suppression affected cancer associated phenotypes in these cell lines.
9
10

11 12 13 14 15 16 **Knockdown of *Nfkb2* exclusively increased the motility of 4T1 cells and this increase** 17 18 **correlated with induction of EMT** 19 20

21 We therefore examined the effects of *Nfkb2* knockdown on the proliferative potential of
22
23 mammary cell lines. Despite the previously described association between p52, cyclin D1 and
24
25 proliferation in transformed cells [25], we did not observe any effects on the growth rate of cells
26
27 (Online Resources 1a-b). We also assessed the consequences of *Nfkb2* knockdown on the colony
28
29 forming potential of these cell lines but no significant differences were observed in the number
30
31 of colony forming units (Online Resources 1a-c). A characteristic of tumour progression to
32
33 malignancy is the phenotypic transition of a sub-population of tumour cells to become more
34
35 motile and invade surrounding tissues. We addressed the role of p100/p52 in the regulation of
36
37 these phenotypes *in vitro* through transwell migration assays. The loss of p100/p52 increased the
38
39 motility of 4T1 cells but had no effect on N202.1A cells (Fig. 2a, Online Resource 2). Motility
40
41 was unaffected in N202.1A cells despite decreases in NF- κ B activity (Fig. 1e), suggesting that
42
43 the alternative NF- κ B pathway is not essential for motility in these cells. As the increase in
44
45 motility of 4T1 cells correlated with an increase in basal NF- κ B activity when *Nfkb2* was
46
47 silenced (Fig. 1e), we wanted to address whether the increase in motility was dependent on NF-
48
49 κ B activity. Treatment of 4T1 *Nfkb2* kd cells with the NF- κ B inhibitor BAY 11-7082
50
51 substantially decreased the motility of these cells to a level that was lower than untreated non-
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 target (NT) control (Fig. 2b). Correspondingly, treatment of 4T1 NT cells with BAY 11-7082
5
6 also decreased motility to a level that was comparable to BAY 11-7082 treated 4T1 Nfkb2 kd
7
8 cells (Fig. 2b). This implied that NF-κB activation was required not only for the enhanced
9
10 motility induced by silencing *Nfkb2* in 4T1 cells, but also the inherent motility of these cells. In
11
12 order to further dissect whether the loss of either p100 or p52 was responsible for the increase in
13
14 motility of 4T1 Nfkb2kd cells, we re-expressed p52 in these cells but this failed to rescue the
15
16 increase in motility due to silencing of *Nfkb2* (Fig. 2c). This suggested that it was the loss of
17
18 p100 that was most likely responsible for the increase in motility of 4T1 Nfkb2kd cells.
19
20
21
22

23
24 Since NF-κB activation is capable of inducing epithelial to mesenchymal transition
25
26 (EMT) and EMT can confer cancer cells with increased migratory properties [33], we were
27
28 interested in determining whether 4T1 Nfkb2kd cells acquired an EMT phenotype. The transcript
29
30 levels of the EMT-inducing transcription factor *Zeb2* were increased 1.6 fold in 4T1 Nfkb2kd
31
32 cells compared to controls (Fig. 2d). Accordingly, we observed a decrease in levels of E-
33
34 cadherin and an increase in levels of N-cadherin relative to NT controls, which is characteristic
35
36 of the EMT process.
37
38
39
40
41
42
43
44
45

46 **The changes in basal NF-κB activity as a result of *Nfkb2* knockdown correlated with the** 47 **changes in CSC properties of 4T1 and N202.1a cells respectively** 48 49

50
51 The NF-κB pathway has been implicated in the regulation of breast CSC maintenance
52
53 and proliferation [21, 22, 29]. In view of that, we wanted to determine whether p100/p52 had a
54
55 role in governing such properties. When we analyzed the levels of putative breast CSC (i.e.
56
57 CD29^{hi}CD61⁺ cells) [34], a significant increase in the proportion of CSCs was observed for 4T1
58
59
60
61
62
63
64
65

cells (Fig. 3a and 3b) and on the other hand, CSC numbers were decreased by more than two fold for N202.1A cells (Fig. 3a and 3c). These changes led us to address CSC phenotypes by employing tumoursphere formation assays in serial passage. In serial passage there was a two fold increase in the proportion of tumourspheres when *Nfkb2* was silenced in 4T1 cells (Fig. 4a) while in N202.1A cells, a significant decrease in tumourspheres was observed following downregulation of *Nfkb2* (Fig. 4a). Consequently, we tested the dependence on NF- κ B activity of respective tumoursphere forming potentials for 4T1 and N202.1A cell lines through inhibition with BAY 11-7082. Notably, BAY 11-7082 treatment abolished the tumoursphere forming potentials of both cell lines to almost undetectable levels (Fig. 4b and 4c). This shows that the inherent tumoursphere forming potential of 4T1 and N202.1A cells upon *Nfkb2* silencing are dependent on NF- κ B signalling. Over-expression of p52 in 4T1 *Nfkb2* kd did not rescue the increased tumoursphere forming potential relative to 4T1 NT cells (Fig. 4d) further suggesting a role for un-processed p100 in 4T1 cell tumorigenesis. However, overexpression of p52 could rescue the decrease in sphere forming efficiency of N202.1A *Nfkb2* kd cells (Fig. 4e), demonstrating a role for p52 in positively regulating tumoursphere forming potential.

Silencing of *Nfkb2* had opposing effects on the tumorigenic potentials of 4T1 and N202.1a cells respectively in vivo

These changes in CSC marker expression and mammosphere forming potential implicate the involvement of p100/p52 in the regulation of tumour initiating properties. Furthermore *in vitro* experiments indicated that 4T1 cells exhibited more aggressive phenotypes when p100/p52 was depleted. To confirm these findings *in vivo* 4T1 NT and *Nfkb2* kd cells were transplanted

1
2
3
4 into the mammary fat pads of recipient mice. 4T1 Nfkb2 kd cells exhibited enhanced growth
5
6 kinetics (Fig. 5a). Although 4T1 Nfkb2 kd tumours were growing relatively faster than 4T1 NT
7
8 tumours, we did not observe any differences in the proliferation or the number of apoptotic cells
9
10 between the two cohorts, as determined by Ki67 and cleaved caspase-3 immuno-staining
11
12 respectively (Fig. 5b-d). This correlated with our findings in vitro (Online Resource 1),
13
14 suggesting that the difference in tumour growth may be due to increased numbers of tumour
15
16 stem/progenitor cells within the 4T1 Nfkb2 kd population.
17
18
19
20
21

22 Concordantly, the decrease in N202.1A CSC activity observed in vitro correlated with a
23
24 decrease in rate of tumour growth of N202.1A xenografts lacking *Nfkb2* (Fig. 5e). Similarly, we
25
26 did not observe any significant difference in the rate of proliferation or apoptosis between the
27
28 N202.1a tumor cohorts as indicated by Ki-67 and cleaved caspase-3 staining (Fig. 5f-h). It is
29
30 possible therefore that the retardation in N202.1a Nfkb2 kd tumor growth was due to decreased
31
32 stem/progenitor cells at the onset of transplant rather than a more general effect on proliferation
33
34 or cell death.
35
36
37
38
39
40
41
42

43 **NMS-873 treatment reduced the sphere forming potential of N202.1A and SKBR3 cells**

44
45

46 Based on the contrasting effects observed in 4T1 and N202.1A cells and the potential role
47
48 for un-processed p100 in these responses, we speculated that inhibition of p100 processing might
49
50 be a more viable therapeutic strategy to avoid the loss of p100 and reduce p52 levels
51
52 concurrently. For that reason, we inhibited an upstream regulator of p100 processing, the
53
54 p97/VCP ATPase [19], using NMS-873, a potent and selective inhibitor of this ATPase. In serial
55
56 passage tumoursphere cultures, we observed a decrease in the sphere forming potential of
57
58
59
60
61
62
63
64
65

1
2
3
4 N202.1A cells after NMS-873 treatment (Fig. 6A), which is consistent with our observations of
5
6 *Nfkb2* silencing in this cell line (Fig. 4A). Importantly, although there is no decrease in the
7
8 sphere forming potential of 4T1 cells in the presence of NMS-873 (Fig. 6A), these cells did not
9
10 exhibit increased sphere forming potential, as observed in 4T1 *Nfkb2* kd cells (Fig. 4A). These
11
12 results also suggest that ErbB2^{HIGH} cells may be more sensitive to p97 inhibition. We expanded
13
14 these observations into two human breast cancer cell lines, MDA-MB-231 (ErbB2^{LOW}) and
15
16 SKBR3 (ErbB2^{HIGH}), and similarly, NMS-873 was only effective in reducing sphere formation in
17
18 SKBR3 cells (Fig. 6B). We also examined the expression of CSC markers, CD24 and CD44
19
20 upon NMS-873 treatment in these cell lines. In MDA-MB-231 cells, no significant changes in
21
22 the CD44⁺CD24⁻ or CD44⁺CD24⁺ populations were observed (Online resource 3). As for
23
24 SKBR3 cells, a CD44⁺CD24⁻ CSC population is not observed in this cell line. However, an
25
26 increase in the CD44⁺CD24⁺ population was apparent in NMS-873 treated SKBR3 cells along
27
28 with a decrease in the CD44⁻CD24⁺ population (Online Resource 3). Interestingly, when we
29
30 analyzed the expression of p97/VCP mRNA levels from TCGA Breast Invasive Carcinoma
31
32 samples using the cbiportal website (<http://cbiportal.org>) [35, 36], the levels of p97 transcript
33
34 was highest in the Her2-enriched breast cancer subtype. Taken together, these findings suggest
35
36 that inhibiting the processing of p100 may be effective to suppress CSC activity in Her2-
37
38 enriched breast tumours.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Discussion

Through silencing of the *Nfkb2* gene in mammary cancer cell lines, we have demonstrated that concomitant loss of the *Nfkb2* gene products, p100 and p52, leads to context dependent effects on the tumorigenicity of these cells. In 4T1 cells, we observed increased NF- κ B activity, induction of EMT, increased motility, CSC properties and tumorigenicity in vivo. Our results are in line with previous studies which have shown the regulation of malignant tumor characteristics and CSC properties by the NF- κ B pathway [21, 33, 37, 38]. We propose that the increase in NF- κ B activity after *Nfkb2* silencing may be due to the loss of p100 which plays more prominent negative regulatory roles in this cell type, suppressing both the canonical and alternative NF- κ B pathways [28]. It is more likely that the consequences of p100 depletion in these cells are mediated through the canonical pathway because reconstitution with p52 did not rescue or aggravate the increased motility and CSC traits in our studies.

On the other hand, we showed that the CSC properties and tumorigenicity of N202.1A cells were positively regulated by p52. Significant decreases in CSC properties were observed in these cells upon *Nfkb2* knockdown and the sphere forming potential could be rescued by reconstitution of p52. These results coincide with a recently published study which implicated the role of the alternative NF- κ B pathway in the regulation of breast CSCs [29]. Furthermore, upstream activators of the alternative NF- κ B pathway such as NIK and IKK α have also been shown to positively regulate breast CSCs [39, 40]. As our results suggest a regulatory role for p100 in limiting CSC activity, the consequences of increased NIK/IKK α activity may be two-fold, losing the inhibitory function of p100 along with increased p52 activity. It is also worth noting that in the study where NIK was implicated in regulation of breast CSCs [39], examination of NIK protein levels through immunohistochemistry revealed that NIK expression

was significantly higher in the Her2⁺ subtype and there was a positive correlation between NIK and Her2 levels. Similarly, in the study where the role of IKK α was examined in several different mouse models of breast cancer, loss of IKK α only affected tumor initiating cells in the MMTV-Neu model [40].

It is possible that not all breast cancers will exhibit a dependence on the alternative NF- κ B pathway for ‘stemness’ and malignancy. Hence, it will be important to identify specific subtypes of breast cancer which will benefit from inhibition of the alternative NF- κ B pathway. At present, it is known that NF- κ B activity in general is elevated in Her2-enriched and basal subtypes of breast cancer [41, 42] but specifically, the alternative NF- κ B pathway has not been associated with a more defined subset of breast cancers. Our data along with studies on upstream regulators of the alternative NF- κ B pathway [39, 40] would indicate that the alternative NF- κ B pathway may have a more important role in Her2-enriched breast tumors. We have also previously shown that Bcl3, a protein which can form complexes with p52 or p50 dimers to activate NF- κ B, specifically plays a role in metastasis of ErbB2 driven tumors [43]. It would be interesting to address whether activation of the alternative NF- κ B pathway is required for the tumor promoting properties of Bcl3.

Although the stratification of breast tumors into molecular subtypes have improved therapeutic outcomes and assisted in administration of specific therapies, current designation of tumor subtypes are based on the bulk of the tumor. Based on the CSC concept, it is possible that multiple breast cancer subtypes (Basal + Luminal) may be present within a tumour [14, 44]. With regards to that, the expression of Her2 has been shown to enrich for a luminal cancer stem cell population [45, 46] and plasticity for this Her2⁺ CSC population has been documented, even in tumors classified as Her2⁻ [47]. Hence it would be interesting to address whether our findings

1
2
3
4 may apply to Her2⁺ CSC subpopulations in breast tumours which are not classified as the Her2
5
6 enriched subtype.
7
8
9

10 Importantly, the context dependent effects we observed in different mammary cancer cell
11 lines would advocate against targeting the *Nfkb2* gene at the genetic level. Accordingly, it would
12 be more appropriate to inhibit the processing of p100 into p52. By employing this therapeutic
13 strategy, p52 transcriptional activity can be repressed and concomitantly, p100 levels can be
14 maintained at higher levels. Since it is possible that p100 can also sequester canonical NF-κB
15 subunits, targeting the processing of p100 could provide additional benefits. We have shown that
16 inhibition of p97 can be effective in limiting CSC activity in a specific subset of breast cancer
17 cells but other potential drug candidates include upstream kinases such as IKKα and NIK. In
18 addition, RANKL inhibitors such as denosumab are already being employed clinically [48] and
19 may prove to be effective in ameliorating certain breast CSCs.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 It appears that after NMS-873 treatment, there was increased CD44 expression in CD24⁺
36 SKBR3 cells (Online Resource 3) which was inversely associated with its sphere forming
37 potential (Figure 6B). Although CD44 positivity is associated with BCSC phenotypes [9], it is
38 possible that only the combined expression pattern of CD44⁺CD24⁻ enriches for BCSCs.
39 Accordingly, previous studies of clinical samples have showed an association between CD44
40 positivity[49-51] and good prognosis but this observations may be convoluted because of the
41 existence of various CD44 isoforms.
42
43
44
45
46
47
48
49
50
51
52

53 In summary, we have shown that the *Nfkb2* gene products p100 and p52, through their
54 roles in the NF-κB pathway, regulate breast CSC properties. We found that p100 may have more
55 significant negative regulatory effects in 4T1^{ERBB2LOW} cells whereas p52 activation was more
56
57
58
59
60
61
62
63
64
65

1
2
3
4 dominant in N202.1A^{ERBB2HIGH} cells. This led to contrasting effects on CSC properties and
5
6
7 tumourigenicity upon *Nfkb2* silencing. The opposing roles of p100 and p52 in the regulation of
8
9
10 breast CSCs gives credence to the inhibition of p100 processing into p52 as a potential strategy
11
12 to target breast CSCs.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgement

We are grateful to Derek Scarborough from the Histology Unit for assistance with histology and processing of tissues and the animal research facility at Cardiff University. We also appreciate the kind feedback and assistance given by Nader Omidvar, Alison Wakefield, Luke Piggott and Joseph Farmer. This work was supported by a Breast Cancer Campaign PhD studentship grant (2007NovPhD06) awarded to R.Clarkson. We would also like to acknowledge the contribution of specimen donors and research groups who have contributed to the TCGA Breast dataset.

Compliance with ethical standards

Conflict of Interest

Authors declare no conflict of interest.

Ethical Standards

The authors declare that the experiments performed in the current publication comply with the current laws of the United Kingdom and all maintenance, breeding and scientific procedures involving animals were carried out according to the guidelines set by the U.K. Home Office Regulations Animals (Scientific Procedures) Act 1986.

References

1. Northern Ireland Cancer Registry (2010) Cancer Incidence and Mortality.
2. Office for National Statistics (2010) Cancer Statistics registrations: Registrations of cancer diagnosed in 2007, England.
3. Welsh Cancer Intelligence and Surveillance Unit (2010) Cancer Incidence in Wales.
4. Ferlay J, et al. (2013) GLOBOCAN 2012 v1.0, Cancer Incidence and mortality worldwide. IARC CancerBase No.11, Lyon, France
5. Perou C, Sorlie T, Eisen M (2000) Molecular portraits of human breast tumours. Nature 406: 747-52
6. Sorlie T, Perou C, Tibshirani R (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. PNAS 98 19): 10869-74
7. Furth J, Kahn M (1937) The transmission of leukemia of mice with a single cell. Am J Cancer 31: 276-82
8. Lapidot T, et al. (1994) A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. Nature 367: 645-8
9. Al-Hajj M, et al. (2003) Prospective identification of tumorigenic breast cancer cells. PNAS 100 7): 3983-8
10. Li X, et al. (2008) Intrinsic resistance of tumorigenic breast cancer cells to chemotherapy. J Natl Cancer Inst 100 9): 672-9
11. Diehn M, et al. (2009) Association of reactive oxygen species levels and radioresistance in cancer stem cells. Nature 458 7239): 780-3
12. Tam WL, et al. (2013) Protein kinase C alpha is a central signaling node and therapeutic target for breast cancer stem cells. Cancer Cell 24 3): 347-64
13. Piggott L, et al. (2011) Suppression of apoptosis inhibitor c-FLIP selectively eliminates breast cancer stem cell activity in response to the anti-cancer agent, TRAIL. Breast Cancer Res 13 5): R88
14. Yeo SK, et al. (2016) Autophagy Differentially Regulates Distinct Breast Cancer Stem-like Cells in Murine Models via EGFR/Stat3 and Tgf β /Smad Signaling. Cancer Res 76 11): 3397-410

15. Baud V, Karin M (2009) Is NF- κ B a good target for cancer therapy? Hopes and pitfalls. *Nature Reviews Drug Discovery* 8: 33-40
16. Perkins N (2007) Integrating cell-signalling pathways with NF- κ B and IKK function. *Nature Reviews Molecular Cell Biology* 8: 49-62
17. Busino L, et al. (2012) Fbxw7 α - and GSK3-mediated degradation of p100 is a pro-survival mechanism in multiple myeloma. *Nat Cell Biol* 14 4): 375-85
18. Fukushima H, et al. (2012) SCF(Fbw7) modulates the NF κ B signaling pathway by targeting NF κ B2 for ubiquitination and destruction. *Cell Rep* 1 5): 434-43
19. Zhang Z, et al. (2015) The Transitional Endoplasmic Reticulum ATPase p97 Regulates the Alternative Nuclear Factor NF- κ B Signaling via Partial Degradation of the NF- κ B Subunit p100. *J Biol Chem* 290 32): 19558-68
20. Curtis C, Shah S, Chin S-F (2012) The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature*
21. Liu M, et al. (2010) The canonical NF-kappaB pathway governs mammary tumorigenesis in transgenic mice and tumor stem cell expansion. *Cancer Research* 70 24): 10464-73
22. Pratt M, et al. (2009) The canonical NF- κ B pathway is required for formation of luminal mammary neoplasias and is activated in the mammary progenitor population. *Oncogene* 8: 2710-22
23. Biswas DK, Iglehart JD (2006) Linkage between EGFR family receptors and nuclear factor kappaB (NF-kappaB) signaling in breast cancer. *J Cell Physiol* 209 3): 645-52
24. Nakshatri H, et al. (1997) Constitutive activation of NF-kappaB during progression of breast cancer to hormone-independent growth. *Mol Cell Biol* 17 7): 3629-39
25. Rocha S, et al. (2003) p53 Represses Cyclin D1 Transcription through Down Regulation of Bcl-3 and Inducing Increased Association of the p52 NF- κ B Subunit with Histone Deacetylase 1. *Molecular and Cellular Biology* 23 13): 4713-27
26. Viatour P, et al. (2003) NF-kappa B2/p100 induces Bcl-2 expression. *Leukemia* 17: 1349-56
27. Cogswell P, et al. (2000) Selective activation of NF-kappa B subunits in human breast cancer: potential roles for NF-kappa B2/p52 and for Bcl-3. *Oncogene* 19 9): 1123-31

- 1
2
3
4 28. Dejardin E, et al. (1995) Highly-expressed p100/p52 (NF- κ B2) sequesters other NF- κ B-related proteins in the cytoplasm of human breast cancer cells. *Oncogene* 11 9): 1835-41
5
6
7
8 29. Kendellen MF, et al. (2014) Canonical and non-canonical NF- κ B signaling promotes
9 breast cancer tumor-initiating cells. *Oncogene* 33 10): 1297-305
10
11
12 30. Aslakson CJ, Miller FR (1992) Selective events in the metastatic process defined by
13 analysis of the sequential dissemination of subpopulations of a mouse mammary tumor. *Cancer*
14 *Res* 52: 1399-405
15
16
17 31. Nanni P, et al. (2000) p185neu protein is required for tumor and anchorage-
18 independent growth, not for cell proliferation of transgenic mammary carcinoma. *Int J Cancer*
19 87: 186-94
20
21
22 32. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
23 quantitative PCR and the 2-delta-deltaCT method. *Methods* 25 4): 402-8
24
25
26 33. Huber MA, et al. (2004) NF- κ B is essential for epithelial-mesenchymal transition and
27 metastasis in a model of breast cancer progression. *J Clin Invest* 114: 569-81
28
29
30 34. Kouros-Mehr H, et al. (2008) GATA-3 links tumor differentiation and dissemination in a
31 luminal breast cancer model. *Cancer Cell* 13 2): 141-52
32
33
34 35. Cerami E, et al. (2012) The cBio cancer genomics portal: an open platform for exploring
35 multidimensional cancer genomics data. *Cancer Discov* 2 5): 401-4
36
37
38 36. Gao J, et al. (2013) Integrative analysis of complex cancer genomics and clinical profiles
39 using the cBioPortal. *Sci Signal* 6 269): p11
40
41
42 37. Cao Y, Luo J, Karin M (2007) I κ B kinase kinase activity is required for self-renewal of
43 ErbB2/Her2-transformed mammary tumor-initiating cells. *PNAS* 104 40): 15852-57
44
45
46 38. Kendellen MF, et al. (2014) Canonical and non-canonical NF- κ B signaling promotes
47 breast cancer tumor-initiating cells. *Oncogene* 33 10): 1297-305
48
49
50 39. Vazquez-Santillan K, et al. (2016) NF- κ B-inducing kinase regulates stem cell
51 phenotype in breast cancer. *Sci Rep* 6: 37340
52
53
54 40. Cao Y, Luo JL, Karin M (2007) I κ B kinase alpha kinase activity is required for self-
55 renewal of ErbB2/Her2-transformed mammary tumor-initiating cells. *Proc Natl Acad Sci U S A*
56 104 40): 15852-7
57
58
59
60
61
62
63
64
65

- 1
2
3
4 41. Yamaguchi N, et al. (2009) Constitutive activation of nuclear factor-kappaB is
5 preferentially involved in the proliferation of basal-like subtype breast cancer cell lines. *Cancer*
6 *Sci* 100 9): 1668-74
7
8
9
10 42. Yamamoto M, et al. (2010) Epigenetic alteration of the NF-κB-inducing kinase (NIK) gene
11 is involved in enhanced NIK expression in basal-like breast cancer. *Cancer Sci* 101 11): 2391-7
12
13
14 43. Wakefield A, et al. (2013) Bcl3 selectively promotes metastasis of ERBB2-driven
15 mammary tumors. *Cancer Res* 73 2): 745-55
16
17
18 44. Liu S, et al. (2014) Breast Cancer Stem Cells Transition between Epithelial and
19 Mesenchymal States Reflective of their Normal Counterparts. *Stem Cell Reports* 2 1): 78-91
20
21
22 45. Ithimakin S, et al. (2013) HER2 drives luminal breast cancer stem cells in the absence of
23 HER2 amplification: implications for efficacy of adjuvant trastuzumab. *Cancer Res* 73 5): 1635-
24 46
25
26
27 46. Korkaya H, et al. (2008) HER2 regulates the mammary stem/progenitor cell population
28 driving tumorigenesis and invasion. *Oncogene* 27 47): 6120-30
29
30
31 47. Jordan NV, et al. (2016) HER2 expression identifies dynamic functional states within
32 circulating breast cancer cells. *Nature* 537 7618): 102-6
33
34
35 48. Drooger JC, et al. (2013) Denosumab in breast cancer treatment. *Eur J Pharmacol* 717 1-
36 3): 12-9
37
38
39 49. Ahmed MA, et al. (2012) A CD44⁻/CD24⁺ phenotype is a poor prognostic marker in early
40 invasive breast cancer. *Breast Cancer Res Treat* 133 3): 979-95
41
42
43 50. Dan T, et al. (2014) CD44 is prognostic for overall survival in the NCI randomized trial on
44 breast conservation with 25 year follow-up. *Breast Cancer Res Treat* 143 1): 11-8
45
46
47 51. Martin TA, Jiang WG (2014) Evaluation of the expression of stem cell markers in human
48 breast cancer reveals a correlation with clinical progression and metastatic disease in ductal
49 carcinoma. *Oncol Rep* 31 1): 262-72
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legends

Figure 1: Silencing of *Nfkb2* alters the basal NF- κ B activity of mammary cell lines. **a** Boxplot of NFKB2 gene expression in tissue samples from TCGA breast dataset acquired through Oncomine software and based upon data generated by the TCGA Research Network: <http://cancergenome.nih.gov/>. Horizontal bars indicate median values, box edges represent 75th and 25th percentile, error bars show 90th percentile and 10th percentile, points show outlier values. **b** Immunoblots of nuclear and cytoplasmic extracts from 4T1 and N202.1A cells showing levels of NF- κ B subunits. **c** Quantification of the ratio of nuclear p52 to nuclear p100 from three independent immunoblots of 4T1 and N202.1A cells. **d** Quantification of the ratio of nuclear p52 to cytoplasmic p52 from three independent immunoblots of 4T1 and N202.1A cells. **e** Bar chart showing relative light units (RLU) from NF- κ B reporter assays. Cells were transfected with NF- κ B luciferase reporter and pcDNA3.1:lacZ as transfection control. 48 hours after transfection, cell lysates were harvested and the ratio of luciferase to lacZ activity measured. Bar charts show basal NF- κ B activity plotted as relative light units (RLU) in the respective cell lines. Data points represent average of at least n=6 and error bars indicate standard error of the mean (SEM). * indicates p<0.05, ** indicates p<0.01 and *** indicates p<0.001.

Figure 2: Silencing of *Nfkb2* increases the motility of 4T1 cells and this increase is associated with EMT in these cells. **a** 4T1 and N202.1a cell lines transduced with non target (NT) and shRNA against *Nfkb2* were plated in Boyden trans-well migration chambers and migration was induced via a serum gradient. Bar charts show the relative number of cells migrated (normalized

1
2
3
4 to 4T1 NT). **b** 4T1 NT and 4T1 Nfkb2 kd cells were plated in Boyden trans-well migration
5
6 chambers in the presence or absence of 1uM BAY 11-7082 and migration was induced via a
7
8 serum gradient. Bar charts show the number of cells migrated. **c** 4T1 cells transduced with non
9
10 target (NT) shRNA, shRNA against Nfkb2 and shRNA against Nfkb2 with p52 over-expression
11
12 were plated in Boyden trans-well migration chambers and migration was induced by a serum
13
14 gradient. Bar charts show number of cells migrated normalized against NT control. **d** qPCR
15
16 analysis of transcripts from 4T1 NT and 4T1 Nfkb2 kd cells. Levels of EMT inducing
17
18 transcription factors were quantified as relative levels of transcripts normalized against NT
19
20 controls using the $\Delta\Delta CT$ method. Data points represent average of at least n=6 and error bars
21
22 indicate standard error of the mean (SEM). * indicates $p<0.05$, where statistical significance
23
24 relative to NT controls were determined by two-tailed t-test.
25
26
27
28
29
30
31
32
33
34

35 **Figure 3:** Effects of silencing *Nfkb2* on the proportion of putative CSCs in mammary cancer cell
36
37 lines. **a** Bar chart showing the percentage of CD29+ CD61+ CSCs in 4T1 and N202.1A cell lines
38
39 transduced with non target (NT) or shRNA against *Nfkb2*. Data points represent average of at
40
41 least n=6 and error bars indicate standard error of the mean (SEM), *** indicates $p<0.001$, where
42
43 statistical significance relative to NT controls were determined by two-tailed t-test.
44
45 Representative dot plots showing percentage of CD29+ CD61+ CSCs in **(b)** 4T1 and **(c)**
46
47 N202.1A cell lines transduced with non target (NT) or shRNA against *Nfkb2*.
48
49
50
51
52
53
54
55

56 **Figure 4:** Effects of silencing *Nfkb2* on the CSC properties of mammary cell lines. **a** 4T1 and
57
58 N202.1A cell lines transduced with non target (NT) or shRNA against *Nfkb2* were plated in non-
59
60
61
62
63
64
65

1
2
3
4 adherent culture plates with mammosphere media. Bar charts show % mammosphere forming
5
6 units (MFUs) in Passage 2 calculated as the number of spheres per cells seeded. **(b)** 4T1 and **(c)**
7
8 N202.1a cell lines in the first passage (P1) with or without 1uM BAY 11-7082 were dissociated
9
10 and re-seeded at an equal density (P2) under non-adherent culture conditions in mammosphere
11
12 media. The number of spheres formed in the presence or absence of BAY 11-7082 respectively
13
14 after 7 days were counted. **(d)** 4T1 and **(e)** N202.1a cells transduced with non target (NT)
15
16 shRNA, shRNA against *Nfkb2* and shRNA against *Nfkb2* with p52 over-expression were plated
17
18 in non-adherent culture plates with mammosphere media and the number of spheres formed after
19
20 7 days in passage 2 was counted. Bar charts show the number of spheres formed. Data points
21
22 represent average of at least n=6 and error bars indicate standard error of the mean (SEM). *
23
24 indicates p<0.05, ** indicates p<0.01, *** indicates p<0.001, n.s. indicates not significant, where
25
26 statistical significance relative to NT controls were determined by two-tailed t-test.
27
28
29
30
31
32
33
34
35
36
37

38 **Figure 5:** Effects of silencing *Nfkb2* on the tumorigenic potential of mammary cancer cells in
39
40 vivo. **a** 10,000 4T1 NT or *Nfkb2* kd cells were transplanted into the right mammary fat pads of
41
42 Balb/C mice respectively. Line graphs show the tumour growth kinetics of respective cell lines. *
43
44 indicates p<0.05, where statistical significance between growth curves was determined by
45
46 ANCOVA test; F=8.370, p=0.018. **b** Representative images from 4T1 NT and *Nfkb2* kd tumours
47
48 immuno-labelled with antibodies against Ki67 and cleaved caspase-3 respectively. Scale bar
49
50 represents 100um. **c** Bar charts showing number of Ki67 positive nuclei per 10x field of view. **d**
51
52 Bar charts showing number of cleaved caspase-3 positive cells per 10x field of view. **e** 50,000
53
54 N202.1A NT or *Nfkb2* kd cells were transplanted into the right mammary fat pads of NOD/SCID
55
56 mice respectively. The resulting tumours were measured and the growth kinetics plotted as line
57
58
59
60
61
62
63
64
65

graphs. Data points represent averages from n=5 animals per cohort and error bars represent standard error of the mean. * indicates $p<0.05$, where statistical significance between growth curves was determined by ANCOVA test; $F=7.754$, $p=0.021$. **f** Representative images from N202.1a NT and *Nfkb2* kd tumours immuno-labelled with antibodies against Ki67 and cleaved caspase-3 respectively. Scale bar represents 100um. **g** Bar charts showing number of Ki67 positive nuclei per 10x field of view. **h** Bar charts showing number of cleaved caspase-3 positive cells per 10x field of view.


Figure 6: NMS-873 inhibits the sphere forming potential of ErbB2 high mammary cancer cells. **a** 4T1 and N202.1A cell lines transduced with non target (NT) or shRNA against *Nfkb2* were plated in non-adherent culture plates with mammosphere media in the absence or presence of 1µM NMS-873. Bar charts show sphere forming efficiency in Passage 2 calculated as the number of spheres per cells seeded. **b** Bar charts showing the sphere forming efficiency of MDA-MB-231 and SKBR3 cells in the absence or presence of 1µM NMS-873. Data points represent average of at least n=6 and error bars indicate standard error of the mean (SEM). ** indicates $p<0.01$, *** indicates $p<0.001$, where statistical significance relative to NT controls were determined by two-tailed t-test. **c** Box plots showing the levels of VCP/p97 mRNA levels in patients from the Breast Invasive Carcinoma TCGA cohort grouped according to PAM50 subtypes. Data was obtained using the cbiportal website and based upon data generated by the TCGA Research Network: <http://cancergenome.nih.gov/>. Statistical significance was determined using ANOVA test; $F=8.69$, $p<0.0001$.


Online Resource 1: Silencing of Nfkb2 does not affect the proliferation and colony forming potential of mammary cancer cell lines. **a** Immunoblot showing the levels of p100 and p52 in 4T1 and N202.1A cells after transduction with Non-Target (NT) or Nfkb2 shRNA. Growth curves showing the number of cells over time for **(b)** 4T1 and **(c)** N202.1A cells. **d** Colony forming assay of 4T1 and N202.1A cells seeded in 6-well plates at a density of 1000 cells/well and quantified after 7 days. Data points represent average of at least n=6 and error bars indicate standard error of the mean (SEM). N.S. indicates a non significant difference, where statistical significance relative to NT controls were determined by two-tailed t-test.


Online Resource 2: Silencing of Nfkb2 with independent shRNAs in 4T1 cells. **a** Immunoblot showing levels of p100 and p52 after knockdown with non target, p52 sh43 or p52 sh44 in 4T1 cells. **b** Bar chart showing number of migrated cells in a 10x field of view for Boyden chamber migration assays of 4T1 non target, p52 sh43 or p52 sh44 cells. **c** Bar chart showing number of spheres formed for mammosphere forming assays of 4T1 non target, p52 sh43 or p52 sh44 cells.

Online Resource 3: Effects of NMS-873 treatment on CSC marker expression in human breast cancer cell lines. **a** Representative dot plots showing expression of CD44⁺ and CD24⁺ in MDA-MB-231 and SKBR3 cell lines treated with vehicle or 1 μ M NMS-873 for 48 hours under normal culture conditions and analyzed by flow cytometry. Bar charts showing the percentage of **b** CD44⁺CD24⁻, **c** CD44⁺CD24⁺, or **d** CD44⁻CD24⁺ populations. Data points represent average of

at least $n=6$ and error bars indicate standard error of the mean (SEM), ** indicates $p<0.01$, where statistical significance relative to vehicle controls were determined by two-tailed t-test.












Figure 4

**Figure 5**

**Figure 6**